



Input of Plant Nutrition

Soluble powder

Reg.: RSCO-188/X/09

CAUTIONS:

In case of ingestion or contact with eyes, take the patient to the doctor and show him the instructive. Store in a cool dry place. Dispose the container appropriately. Do not use the container again. Keep out of reach of children.

COMPATIBILITY: The available research does not indicate incompatibility with agrochemicals frequently used. Cut costs by applying ENERPLANT® with mixtures of agrochemicals used for other purposes. Except herbicides.

GUARANTEE:

The company guarantees that the composition of this product fulfills the specifications of our label. Due to its final use is out of our control, the buyer assumes total responsibility for its handling.

Always calibrate your application equipment.

WARNING, READ THE LABEL BEFORE USING THE PRODUCT.

PORCENTUAL COMPOSITION:

Active Ingredient:

Oligosaccharides.....0.01%

Inert Ingredients:

Dextrose79.90%

Maltodextrin.....19.99%

Vegetal Coloring.....0.10%

Total:.....100.00%

NON TOXIC PRODUCT

INSTRUCTIONS: Premixes with water (1 gallon) before incorporate into the mixing tank. We recommend using at least 200 L (52.83 gal) of water to cover one hectare of crop. Or 80 L (21.13 gal) to cover one acre.



Certified for organic agricultura by BCS Oko.



For organic production OMRI.org

DOSAGE AND FREQUENCY OF APPLICATION PER CROP

Crop	Dosage per cycle/ha	Dosage: gr/ha	Application	Moment of application	Via
RICE	1 cycle of 80 gr per hectare in 2 applications	40 gr/ha	1st.	15-25 days after of emerge.	Foliage
		40 gr/ha	2nd.	15-25 days after of the 1st.	Foliage
COTTON	1 cycle of 60 gr per hectare in 3 applications	20 gr/ha	1st.	15-25 days after of germinate.	Foliage
		20 gr/ha	2nd.	15-25 days after of the 1st.	Foliage
		20 gr/ha	3rd.	15-25 days after of the 2nd.	Foliage
BANANA	1 cycle of 400 gr per hectare in 10 applications	40 gr/ha	1 – 10	1 application every month, starting to the 6 months after of transplant.	Foliage
COFFEE	1 cycle of 60 gr per hectare in 3 applications	20 gr/ha	1st.	At starting bloom.	Foliage
		20 gr/ha	2nd.	In pollination	Foliage
		20 gr/ha	3rd.	25-30 days after of the 2nd.	Foliage
SUGAR CANE	1 cycle of 80 gr per hectare in 2 applications	40 gr/ha	1st.	30-35 days after of sprout.	Foliage
		40 gr/ha	2nd.	30-35 days after of the 1st.	Foliage
GRASSES AND FORAGE	1 cycle of 80 gr per hectare in 2 applications	40 gr/ha	1st.	7-10 days after of cut.	Foliage
		40 gr/ha	2nd.	7-10 days after of the 1st.	Foliage
BEAN, SOY, CHICKPEA AND PEA	1 cycle of 60 gr per hectare in 3 applications	20 gr/ha	1st.	10-12 days after of emerge.	Foliage
		20 gr/ha	2nd.	10-12 days after of the 1st.	Foliage
		20 gr/ha	3rd.	10-12 days after of the 2nd.	Foliage
FRUIT TREES	1 cycle of 120 gr per hectare in 3 applications	40 gr/ha	1st.	At starting bloom.	Foliage
		40 gr/ha	2nd.	In pollination	Foliage
		40 gr/ha	3rd.	30 days after of the 2nd.	Foliage
CORN AND SORGHUM	1 cycle of 80 gr per hectare in 2 applications	40 gr/ha	1st.	15-25 days after of sprout.	Foliage
		40 gr/ha	2nd.	25-30 days after of the 1st.	Foliage
WHEAT, BARLEY AND OAT	1 cycle of 80 gr per hectare in 2 applications	40 gr/ha	1st.	15-20 days after of sprout.	Foliage
		40 gr/ha	2nd.	10-15 days after of the 1st.	Foliage
OIL PALM IN PRODUCTION	2 cycles of 240 gr per hectare in 6 applications	40 gr/ha	1st.	30 days before of the fertilization.	Foliage
		40 gr/ha	2nd.	25-30 days after of the 1st.	Foliage
		40 gr/ha	3rd.	25-30 days after of the 2nd.	Foliage
		40 gr/ha	4th.	6 months after of the 1st.	Foliage
		40 gr/ha	5th.	6 months after of the 4th.	Foliage
		40 gr/ha	6th.	6 months after of the 5a.	Foliage
VEGETABLES (SOWING OF SEEDS)	1 cycle of 60 gr per hectare in 3 applications	20 gr/ha	1st.	25-35 days after of germinate.	Foliage
		20 gr/ha	2nd.	15-20 days after of the 1st.	Foliage
		20 gr/ha	3rd.	15-20 days after of the 2nd.	Foliage
VEGETABLES (SOWING OF SEEDLINGS)	1 cycle of 60 gr per hectare in 3 applications	20 gr/ha	1st.	15-20 days after of transplant.	Foliage
		20 gr/ha	2nd.	15-20 days after of the 1st.	Foliage
		20 gr/ha	3rd.	15-20 days after of the 2nd.	Foliage

Crop	Dosage per cycle/ha	Dosage: gr/ha	Application	Moment of application	Vía
POTATO	1 cycle of 60 gr per hectare in 3 applications	20 gr/ha 20 gr/ha 20 gr/ha	1st. 2nd. 3rd.	15-20 days after emerge. 25-30 days after of the 1st. 25-30 days after of the 2nd.	Foliage Foliage Foliage
PAPAYA	1 cycle of 200 gr per hectare in 10 applications	20 gr/ha	1 – 10	1 application every month, starting a the 6 months after of the transplant.	Foliage
PEPPERS	1 cycle of 60 gr per hectare in 3 applications	20 gr/ha 20 gr/ha 20 gr/ha	1st. 2nd. 3rd.	15-20 days after of transplant. 15-20 days after of the 1st. 15-20 days after of the 2nd.	Foliage Foliage Foliage
PINEAPPLE	1 cycle of 60 gr per hectare in 3 applications	20 gr/ha 20 gr/ha 20 gr/ha	1st. 2nd. 3rd.	25-30 days before of the floral induction. 25-30 days after of the floral induction. 15-20 days after of the 2nd.	Foliage Foliage Foliage
TOBACCO	1 cycle of 120 gr per hectare in 3 applications	40 gr/ha 40 gr/ha 40 gr/ha	1st. 2nd. 3rd.	20-25 days after of transplant. 15-20 days after of the 1st. 15-20 days after of the 2nd.	Foliage Foliage Foliage
TOMATO	1 cycle of 60 gr per hectare in 3 applications	20 gr/ha 20 gr/ha 20 gr/ha	1st. 2nd. 3rd.	15-20 days after of transplant. 15-20 days after of the 1st. 15-20 days after of the 2nd.	Foliage Foliage Foliage
ROSES AND PERENNIAL FLOWERS	1 cycle of 60 gr per hectare in 3 applications	20 gr/ha 20 gr/ha 20 gr/ha	1st. 2nd. 3rd.	10-15 days after of the pruning. 10-12 days after of the 1st. Before of starting buttoning	Foliage Foliage Foliage
WATERMELON, MELON, PUMPKIN AND CUCUMBER (SEED)	1 cycle of 60 gr per hectare in 3 applications	20 gr/ha 20 gr/ha 20 gr/ha	1st. 2nd. 3rd.	15-20 days after emerge. 10-15 days after of the 1st. 10-15 days after of the 2nd.	Foliage Foliage Foliage
WATERMELON, MELON, PUMPKIN AND CUCUMBER (SEEDLING)	1 cycle of 60 gr per hectare in 3 applications	20 gr/ha 20 gr/ha 20 gr/ha	1st. 2nd. 3rd.	10-15 days after of transplant. 10-15 days after of the 1st. 10-15 days after of the 2nd.	Foliage Foliage Foliage
CARROT	1 cycle of 60 gr per hectare in 3 applications	20 gr/ha 20 gr/ha 20 gr/ha	1st. 2nd. 3rd.	10-15 days after of germinate. 10-15 days after of the 1st. 10-15 days after of the 2nd.	Foliage Foliage Foliage

✦ **RECOMENDATIONS FOR THE APPLICATION IN SEEDS (OPTIONAL).**

1. Prepare the necessary amount of seed according to the area to be planted, based on the amount of seed that is planted for each hectare of crop area.
2. 20 grams of **Enerplant®** are diluted in a volume of water of 500 ml (or more if necessary) to moisten the amount of seed corresponding to 1 ha.
3. The seed is sprinkled with the **Enerplant®** solution and a blanket or drum can be used to constantly move the seed for about 5 minutes until the seed is aerated and sowing can be done immediately.
4. **Enerplant®** can be mixed with any agrochemical or biological product to treat the seed.