



**Vegetal Nutrition**

*Concentrated liquid*

Reg.: RSCO-0492/X/99

Always calibrate your application equipment.

**PORCENTUAL COMPOSITION:** Weight percentage

**Active Ingredient:**

Oligosaccharides.....	0.01%
Conservator: Ethylic alcohol.....	49.99%
Vehicle: Water.....	49.99%
Vegetal Coloring.....	0.01%
Total:.....	100.00%

**CAUTIONS:**  
In case of ingestion or contact with eyes, take the patient to the doctor and show him the instructive. Store in a cool dry place. Dispose the container appropriately. Do not use the container again. Keep out of reach of children.

**WARNING, READ THE LABEL BEFORE USING THE PRODUCT.**

**NON TOXIC PRODUCT**

**INSTRUCTIONS:** Premixes with water (1 gallon) before incorporate into the mixing tank. We recommend using at least 200 L (52.83 gal) of water to cover one hectare of crop. Or 80 L (21.13 gal) to cover one acre.

**COMPATIBILITY:** The available research does not indicate incompatibility with agrochemicals frequently used. Cut costs by applying ENERPLANT® with mixtures of agrochemicals used for other purposes. Except herbicides.

**GUARANTEE:**  
The company guarantees that the composition of this product fulfills the specifications of our label. Due to its final use is out of our control, the buyer assumes total responsibility for its handling.



Certified for organic agricultura by BCS Oko.



For organic production OMRI.org

**DOSAGE AND FREQUENCY OF APPLICATION PER CROP**

<b>Crop</b>	<b>Dosage per cycle/ha</b>	<b>Dosage: ml/ha</b>	<b>Application</b>	<b>Moment of application</b>	<b>Via</b>
<b>RICE</b>	1 cycle of 5.2 ml per hectare in 2 applications	2.6 ml/ha	1st.	15-25 days after of emerge.	Foliage
		2.6 ml/ha	2nd.	15-25 days after of the 1st.	Foliage
<b>COTTON</b>	1 cycle of 3.9 ml per hectare in 3 applications	1.3 ml/ha	1st.	15-25 days after of germinate.	Foliage
		1.3 ml/ha	2nd.	15-25 days after of the 1st.	Foliage
		1.3 ml/ha	3rd.	15-25 days after of the 2nd.	Foliage
<b>BANANA</b>	1 cycle of 26 ml per hectare in 10 applications	2.6 ml/ha	1 – 10	1 application every month, starting to the 6 months after of transplant.	Foliage
<b>COFFEE</b>	1 cycle of 3.9 ml per hectare in 3 applications	1.3 ml/ha	1st.	At starting bloom.	Foliage
		1.3 ml/ha	2nd.	In pollination	Foliage
		1.3 ml/ha	3rd.	25-30 days after of the 2nd.	Foliage
<b>SUGAR CANE</b>	1 cycle of 5.2 ml per hectare in 2 applications	2.6 ml/ha	1st.	30-35 days after of sprout.	Foliage
		2.6 ml/ha	2nd.	30-35 days after of the 1st.	Foliage
<b>GRASSES AND FORAGE</b>	1 cycle of 5.2 ml per hectare in 2 applications	2.6 ml/ha	1st.	7-10 days after of cut.	Foliage
		2.6 ml/ha	2nd.	7-10 days after of the 1st.	Foliage
<b>BEAN, SOY, CHICKPEA AND PEA</b>	1 cycle of 3.9 ml per hectare in 3 applications	1.3 ml/ha	1st.	10-12 days after of emerge.	Foliage
		1.3 ml/ha	2nd.	10-12 days after of the 1st.	Foliage
		1.3 ml/ha	3rd.	10-12 days after of the 2nd.	Foliage
<b>FRUIT TREES</b>	1 cycle of 7.8 ml per hectare in 3 applications	2.6 ml/ha	1st.	At starting bloom.	Foliage
		2.6 ml/ha	2nd.	In pollination	Foliage
		2.6 ml/ha	3rd.	30 days after of the 2nd.	Foliage
<b>CORN AND SORGHUM</b>	1 cycle of 5.2 ml per hectare in 2 applications	2.6 ml/ha	1st.	15-25 days after of sprout.	Foliage
		2.6 ml/ha	2nd.	25-30 days after of the 1st.	Foliage
<b>WHEAT, BARLEY AND OAT</b>	1 cycle of 5.2 ml per hectare in 2 applications	2.6 ml/ha	1st.	15-20 days after of sprout.	Foliage
		2.6 ml/ha	2nd.	10-15 days after of the 1st.	Foliage
<b>OIL PALM IN PRODUCTION</b>	2 cycles of 15.6 ml per hectare in 6 applications	2.6 ml/ha	1st.	30 days before of the fertilization.	Foliage
		2.6 ml/ha	2nd.	25-30 days after of the 1st.	Foliage
		2.6 ml/ha	3rd.	25-30 days after of the 2nd.	Foliage
		2.6 ml/ha	4th.	6 months after of the 1st.	Foliage
		2.6 ml/ha	5th.	6 months after of the 4th.	Foliage
		2.6 ml/ha	6th.	6 months after of the 5a.	Foliage
<b>VEGETABLES (SOWING OF SEEDS)</b>	1 cycle of 3.9 ml per hectare in 3 applications	1.3 ml/ha	1st.	25-35 days after of germinate.	Foliage
		1.3 ml/ha	2nd.	15-20 days after of the 1st.	Foliage
		1.3 ml/ha	3rd.	15-20 days after of the 2nd.	Foliage
<b>VEGETABLES (SOWING OF SEEDLINGS)</b>	1 cycle of 3.9 ml per hectare in 3 applications	1.3 ml/ha	1st.	15-20 days after of transplant.	Foliage
		1.3 ml/ha	2nd.	15-20 days after of the 1st.	Foliage
		1.3 ml/ha	3rd.	15-20 days after of the 2nd.	Foliage

<b>Crop</b>	<b>Dosage per cycle/ha</b>	<b>Dosage: gr/ha</b>	<b>Application</b>	<b>Moment of application</b>	<b>Via</b>
<b>POTATO</b>	1 cycle of 3.9 ml per hectare in 3 applications	1.3 ml/ha 1.3 ml/ha 1.3 ml/ha	1st. 2nd. 3rd.	15-20 days after emerge. 25-30 days after of the 1st. 25-30 days after of the 2nd.	Foliage Foliage Foliage
<b>PAPAYA</b>	1 cycle of 13 ml per hectare in 10 applications	1.3 ml/ha	1 – 10	1 application every month, starting a the 6 months after of the transplant.	Foliage
<b>PEPPERS</b>	1 cycle of 3.9 ml per hectare in 3 applications	1.3 ml/ha 1.3 ml/ha 1.3 ml/ha	1st. 2nd. 3rd.	15-20 days after of transplant. 15-20 days after of the 1st. 15-20 days after of the 2nd.	Foliage Foliage Foliage
<b>PINEAPPLE</b>	1 cycle of 3.9 ml per hectare in 3 applications	1.3 ml/ha 1.3 ml/ha 1.3 ml/ha	1st. 2nd. 3rd.	25-30 days before of the floral induction. 25-30 days after of the floral induction. 15-20 days after of the 2nd.	Foliage Foliage Foliage
<b>TOBACCO</b>	1 cycle of 5.8 ml per hectare in 2 applications	2.6 ml/ha 2.6 ml/ha	1st. 2nd.	20 days after of transplant. 15 days after of the 1st.	Foliage Foliage
<b>TOMATO</b>	1 cycle of 3.9 ml per hectare in 3 applications	1.3 ml/ha 1.3 ml/ha 1.3 ml/ha	1st. 2nd. 3rd.	15-20 days after of transplant. 15-20 days after of the 1st. 15-20 days after of the 2nd.	Foliage Foliage Foliage
<b>ROSES AND PERENNIAL FLOWERS</b>	1 cycle of 3.9 ml per hectare in 3 applications	1.3 ml/ha 1.3 ml/ha 1.3 ml/ha	1st. 2nd. 3rd.	10-15 days after of the pruning. 10-12 days after of the 1st. Before of starting buttoning	Foliage Foliage Foliage
<b>WATERMELON, MELON, PUMPKIN AND CUCUMBER (SEED)</b>	1 cycle of 3.9 ml per hectare in 3 applications	1.3 ml/ha 1.3 ml/ha 1.3 ml/ha	1st. 2nd. 3rd.	15-20 days after emerge. 10-15 days after of the 1st. 10-15 days after of the 2nd.	Foliage Foliage Foliage
<b>WATERMELON, MELON, PUMPKIN AND CUCUMBER (SEEDLING)</b>	1 cycle of 3.9 ml per hectare in 3 applications	1.3 ml/ha 1.3 ml/ha 1.3 ml/ha	1st. 2nd. 3rd.	10-15 days after of transplant. 10-15 days after of the 1st. 10-15 days after of the 2nd.	Foliage Foliage Foliage
<b>CARROT</b>	1 cycle of 3.9 ml per hectare in 3 applications	1.3 ml/ha 1.3 ml/ha 1.3 ml/ha	1st. 2nd. 3rd.	10-15 days after of germinate. 10-15 days after of the 1st. 10-15 days after of the 2nd.	Foliage Foliage Foliage

✦ **RECOMENDATIONS FOR THE APPLICATION IN SEEDS (OPTIONAL).**

1. Prepare the necessary amount of seed according to the area to be planted, based on the amount of seed that is planted for each hectare of crop area.
2. 1.3 ml of **Enerplant®** are diluted in a volume of water of 500 ml (or more if necessary) to moisten the amount of seed corresponding to 1 ha.
3. The seed is sprinkled with the **Enerplant®** solution and a blanket or drum can be used to constantly move the seed for about 5 minutes until the seed is aerated and sowing can be done immediately.
4. **Enerplant®** can be mixed with any agrochemical or biological product to treat the seed.

